



Year: 2019

Evaluation of Lightmix Mycoplasma macrolide assay for detection of macrolide-resistant Mycoplasma pneumoniae in pneumonia patients

Wagner, K ; Imkamp, F ; Pires, V P ; Keller, P M

Abstract: OBJECTIVES Rapid detection of macrolide resistance-associated mutations in Mycoplasma pneumoniae is crucial for effective antimicrobial treatment. We evaluated the Lightmix Mycoplasma macrolide assay for the detection of point mutations at nucleotide positions 2063 and 2064 in the 23S ribosomal RNA (rRNA) gene of M. pneumoniae that confer macrolide resistance. METHODS Samples from 3438 patients with a respiratory tract infection were analysed by M. pneumoniae real-time PCR, and 208 (6%) of them were tested positive. In this retrospective study, 163 M. pneumoniae real-time PCR-positive samples were analysed by the Lightmix assay, and results were compared to targeted 23S rRNA sequencing. RESULTS Macrolide-resistant M. pneumoniae were found in 15 (9%) of 163 retrospectively analysed samples. The Lightmix assay showed a sensitivity of 100% (95% confidence interval, 78.2-100) and a specificity of 100% (95% confidence interval, 97.5-100) as the detected M. pneumoniae genotype (148 wild type and 15 non-wild type) was confirmed by 23S rRNA sequencing in all samples. CONCLUSIONS The Lightmix assay is an easy-to-use and accurate molecular test that allows rapid determination of macrolide resistance in M. pneumoniae.

DOI: <https://doi.org/10.1016/j.cmi.2018.10.006>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-168528>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.

Originally published at:

Wagner, K; Imkamp, F; Pires, V P; Keller, P M (2019). Evaluation of Lightmix Mycoplasma macrolide assay for detection of macrolide-resistant Mycoplasma pneumoniae in pneumonia patients. Clinical Microbiology and Infection, 25(3):383.e5-383.e7.

DOI: <https://doi.org/10.1016/j.cmi.2018.10.006>



Research note

Evaluation of Lightmix *Mycoplasma* macrolide assay for detection of macrolide-resistant *Mycoplasma pneumoniae* in pneumonia patientsK. Wagner^{*}, F. Imkamp, V.P. Pires, P.M. Keller

Institute of Medical Microbiology, University of Zurich, Zurich, Switzerland

ARTICLE INFO

Article history:

Received 27 July 2018

Received in revised form

8 October 2018

Accepted 12 October 2018

Available online 26 October 2018

Editor: F. Allerberger

Keywords:

Antibiotic resistance

Macrolide

Molecular detection

Mycoplasma pneumoniae

Pneumonia

RT-PCR

Switzerland

ABSTRACT

Objectives: Rapid detection of macrolide resistance—associated mutations in *Mycoplasma pneumoniae* is crucial for effective antimicrobial treatment. We evaluated the Lightmix *Mycoplasma* macrolide assay for the detection of point mutations at nucleotide positions 2063 and 2064 in the 23S ribosomal RNA (rRNA) gene of *M. pneumoniae* that confer macrolide resistance.

Methods: Samples from 3438 patients with a respiratory tract infection were analysed by *M. pneumoniae* real-time PCR, and 208 (6%) of them were tested positive. In this retrospective study, 163 *M. pneumoniae* real-time PCR—positive samples were analysed by the Lightmix assay, and results were compared to targeted 23S rRNA sequencing.

Results: Macrolide-resistant *M. pneumoniae* were found in 15 (9%) of 163 retrospectively analysed samples. The Lightmix assay showed a sensitivity of 100% (95% confidence interval, 78.2–100) and a specificity of 100% (95% confidence interval, 97.5–100) as the detected *M. pneumoniae* genotype (148 wild type and 15 non-wild type) was confirmed by 23S rRNA sequencing in all samples.

Conclusions: The Lightmix assay is an easy-to-use and accurate molecular test that allows rapid determination of macrolide resistance in *M. pneumoniae*. **K. Wagner, Clin Microbiol Infect 2019;25:383.e5–383.e7**

© 2018 The Authors. Published by Elsevier Ltd on behalf of European Society of Clinical Microbiology and Infectious Diseases. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Mycoplasma pneumoniae is a common pathogen causing respiratory tract infections, particularly in children and young adults [1]. Macrolides are used as empiric first-line treatment of *M. pneumoniae* infections. However, their extensive use led to the rapid worldwide emergence of clinically macrolide-resistant *M. pneumoniae* strains [2,3]. The main mechanism of macrolide resistance in *M. pneumoniae* is nucleotide substitutions at positions A2063 and A2064 in domain V of the 23S ribosomal RNA (rRNA) gene (*M. pneumoniae* numbering; corresponds to nucleotide positions 2058 and 2059 in *Escherichia coli*) [4].

The aim of this study was to evaluate the reliability and accuracy of the commercially available molecular Lightmix assay to detect macrolide-resistant *M. pneumoniae* in respiratory specimens of patients with pneumonia.

Methods

Clinical specimens and *M. pneumoniae* real-time PCR

All respiratory specimens ($N = 3438$) used in this study were sent to the Institute of Medical Microbiology, University of Zurich, between January 2014 and December 2017 for the detection of respiratory bacterial pathogens, including *M. pneumoniae*, from secondary or tertiary care facilities, and were recovered from patients with respiratory tract infection symptoms. Isolation of chromosomal DNA and *M. pneumoniae* real-time PCR (RT-PCR) detection (henceforward termed *M. pneumoniae* RT-PCR) was performed on all 3438 respiratory specimens as described previously [5–7].

Lightmix *Mycoplasma* macrolide assay

Chromosomal DNA of respiratory specimens was stored at -20°C . Before conducting this retrospective study, the DNA concentration of all *M. pneumoniae* RT-PCR—positive samples was determined using the Qubit dsDNA HS kit (Thermo Fischer

^{*} Corresponding author: K. Wagner, Division of Infection Diagnostics, Department of Biomedicine, University of Basel, Petersplatz 10, 4051 Basel, Switzerland. E-mail address: karoline.wagner@usb.ch (K. Wagner).

Scientific, Schlieren, Switzerland). Samples with a DNA concentration of >0.5 ng/μL were analysed by the Lightmix *Mycoplasma* macrolide assay (TIBMolbiol, Berlin, Germany) (Supplementary Fig. S1). The Lightmix assay consists of a RT-PCR amplifying a 117 bp fragment of the 23S rRNA gene and subsequent melting curve analysis. The assay was performed on a LightCycler 480-II instrument (Roche, Rotkreuz, Switzerland). A melting temperature (T_m) greater 66°C in the Lightmix assay indicated a wild-type genotype, while a T_m lower than 62°C indicated the presence of a point mutation at position A2063 or 2064 in the amplified 23S rRNA gene, rendering the detected *M. pneumoniae* macrolide resistant. To verify the genotype detected by the Lightmix assay, all 163 23S rRNA amplicons were sequenced and analysed with respect to resistance-conferring point mutations.

23S rRNA RT-PCR amplicon sequencing

The 23S rRNA RT-PCR products were cleaned using AmpPure XP beads (Beckman Coulter, Nyon, Switzerland), and library preparation was done with the QIAseq FX DNA library preparation kit (Qiagen, Hilden, Germany) following the producers' recommendations except omitting the fragmentation step. Sequencing library quality and size distribution was analysed on a fragment analyser (Advanced Analytical Technologies, Heidelberg, Germany) using the Fragment Analyzer 474 HS NGS kit. Sequencing libraries were pooled in equimolar concentrations and paired-end sequenced (2 × 150 bp) on an Illumina MiSeq platform (Illumina, San Diego, CA, USA). Reads of the 23S rRNA gene sequences of the *M. pneumoniae* isolates were mapped to the 23S rRNA gene of *M. pneumoniae* strain M129 using CLC Genomic Workbench software (Qiagen).

Data analysis

A chi-square test of independence was performed to examine the associations among age, proportion of *M. pneumoniae* infections and macrolide resistance. All statistical data analysis and data visualization were performed using R software [8].

Ethics statement

This retrospective study was conducted in accordance with the Declaration of Helsinki and national and institutional standards. The act on medical research involving human subjects does not apply to this study. The study was approved by the ethics committee of the canton of Zurich, Switzerland (Req-2018-00473).

Results

Overall, 208 of the 3438 respiratory specimens were tested positive by *M. pneumoniae* RT-PCR. In 45 of 208 *M. pneumoniae*-positive samples, chromosomal DNA was degraded as a result of prolonged freezing (<0.5 ng/μL DNA; $n = 7$), or samples were no longer available for retrospective testing ($n = 38$) via the Lightmix assay. Of the

remaining 163 samples, 148 (91%) showed a wild-type genotype in the Lightmix assay, while 15 (9%) displayed a mutation at nucleotide position A2063 or A2064 in the 23S rRNA gene conferring resistance to macrolides (Table 1, Supplementary Fig. S1).

Isolates classified as *M. pneumoniae* wild type in the Lightmix assay did not show a mutation in the domain V of the 23S rRNA gene, as verified by 23S rRNA gene amplicon sequencing. For all samples that showed a shifted melting temperature ($T_m < 62^\circ\text{C}$; $n = 15$) in the Lightmix assay, sequencing confirmed a mutation either at nucleotide position 2063 or 2064. We found nine A2063G mutations, five A2064G mutations and one A2063C mutation (Table 1). This resulted in a sensitivity of 100% (95% confidence interval, 78.2–100) and a specificity of 100% (95% confidence interval, 97.5–100) of the Lightmix assay.

Interestingly, in our study, we found that the median age of patients with a *M. pneumoniae* infection was 15 years, while the median age of patients with macrolide-resistant *M. pneumoniae* infections was 47 years (Supplementary Fig. S2). There was a clear association between the age of patients and *M. pneumoniae* infection ($\chi^2 (5, n = 3217) = 328.0, p < 0.001$) with the highest proportion of *M. pneumoniae* infections in patients aged 0 to 20 years (Supplementary Fig. S3(A)). Interestingly, among patients with *M. pneumoniae* infections, younger patients showed infections with macrolide-susceptible *M. pneumoniae*, but macrolide-resistant *M. pneumoniae* infections were more prevalent in patients aged >31 years ($\chi^2 (5, n = 211) = 12.1, p < 0.05$), with the strongest association in patients aged >50 years (Supplementary Fig. S3(B)).

Discussion

In this retrospective study, the Lightmix *Mycoplasma* macrolide assay was evaluated on respiratory specimens of pneumonia patients and showed high specificity and sensitivity for the detection of point mutations in the 23S rRNA gene conferring macrolide resistance in *M. pneumoniae*.

In this study, a macrolide resistance rate of 9% was found in *M. pneumoniae* when analysing samples from patients treated at hospitals in Zurich, Switzerland. Prevalence of macrolide resistance in *M. pneumoniae* differs in Europe and has been previously reported as being 0 to 3% in Germany, the Netherlands and Switzerland [9–11], 9.8% in France [12] and 26% in Italy [13]. Differences in observed resistance rates in this study compared to previous studies conducted in Switzerland [9] may be due to two main factors. Firstly, the respiratory specimens analysed in this study are mainly obtained from patients treated at secondary or tertiary care facilities. Therefore, the patient population may be selected in terms of progression of disease, severity of pneumonia and proportion of persistent or recurrent *M. pneumoniae* infections; further, patients are often already receiving antibiotic therapy when samples are retrieved for diagnostic procedures. Secondly, the age distribution of the patient population is very broad (1 to 98 years old), and therefore resistance rates may differ from studies that primarily focused on *M. pneumoniae* infection and macrolide resistance in children and young adults [9]. This notion is supported by the fact that we found a

Table 1
Retrospective analysis of respiratory specimens by Lightmix assay and confirmation of detected *Mycoplasma pneumoniae* genotype by 23S rRNA amplicon sequencing

<i>M. pneumoniae</i> by RT-PCR		Genotype not determined	Retrospective analysis by Lightmix assay		Genotype determined by 23S rRNA sequencing	
Negative	Positive		WT	Non-WT	WT	Non-WT
3230 (94%)	208 (6%)	45	148 (91%)	15 (9%)	148	A2063C (1 ×), A2063G (9 ×), A2064G (5 ×)

rRNA, ribosomal RNA; RT-PCR, real-time PCR; WT, wild type.

higher proportion of macrolide-resistant *M. pneumoniae* infections in patients aged >31 years. This indicates that macrolide resistance may be associated with age and possibly long-term antibiotic consumption, as macrolides are among the most commonly prescribed antibiotics in Switzerland [14].

This study has some limitations. It was designed as a single-centre laboratory-based, retrospective evaluation study with extracted DNA from clinical respiratory specimens. We found a high macrolide resistance rate in *M. pneumoniae* in the investigated patient population. This would need to be further addressed in future studies, including patients from primary care facilities. Thus, continuous surveillance is crucial to monitor macrolide resistance in *M. pneumoniae* and to further investigate its epidemiology in Switzerland.

Acknowledgements

We thank the technicians of the Institute of Medical Microbiology for expert help and assistance; and we thank the Institute of Medical Microbiology, University of Zurich, for continuous support.

Transparency Declaration

Lightmix kits for the analysis of 100 samples were donated by TIB Molbiol, Berlin, Germany. For the analysis of the remaining samples, Lightmix kits were purchased from TIB Molbiol by the Institute of Medical Microbiology, University of Zurich. All authors report no conflicts of interest relevant to this article.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.cmi.2018.10.006>.

References

- [1] Waites KB, Xiao L, Liu Y, Balish MF, Atkinson TP. *Mycoplasma pneumoniae* from the respiratory tract and beyond. Clin Microbiol Rev 2017;30:747–809.
- [2] Morozumi M, Takahashi T, Ubukata K. Macrolide-resistant *Mycoplasma pneumoniae*: characteristics of isolates and clinical aspects of community-acquired pneumonia. J Infect Chemother 2010;16:78–86.
- [3] Pereyre S, Goret J, Bébér C. *Mycoplasma pneumoniae*: current knowledge on macrolide resistance and treatment. Front Microbiol 2016;7:974.
- [4] Liu X, Jiang Y, Chen X, Li J, Shi D, Xin D. Drug resistance mechanisms of *Mycoplasma pneumoniae* to macrolide antibiotics. Biomed Res Int 2014;2014.
- [5] Nadala D, Bossart W, Zucol F, Steiner F, Berger C, Lips U, et al. Community-acquired pneumonia in children due to *Mycoplasma pneumoniae*: diagnostic performance of a seminested 16S rDNA-PCR. Diagn Microbiol Infect Dis 2001;39:15–9.
- [6] Hardegger D, Nadal D, Bossart W, Altwegg M, Dutly F. Rapid detection of *Mycoplasma pneumoniae* in clinical samples by real-time PCR. J Microbiol Methods 2000;41:45–51.
- [7] Wagner K, Springer B, Imkamp F, Opota O, Greub G, Keller PM. Detection of respiratory bacterial pathogens causing atypical pneumonia by multiplex Lightmix® RT-PCR. Int J Med Microbiol 2018;308:317–23.
- [8] R Development Core Team. R: a language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing; 2013.
- [9] Sauteur PM, Bleisch B, Voit A, Maurer FP, Relly C, Berger C, et al. Survey of macrolide-resistant *Mycoplasma pneumoniae* in children with community-acquired pneumonia in Switzerland. Swiss Med Wkly 2014;144:w14041.
- [10] Dumke R, Schnee C, Pletz MW, Rupp J, Jacobs E, Sachse K, et al. *Mycoplasma pneumoniae* and *Chlamydia* spp. infection in community-acquired pneumonia, Germany, 2011–2012. Emerg Infect Dis 2015;21:426.
- [11] Spuesens EB, Meijer A, Bierschenk D, Hoogenboezem T, Donker GA, Hartwig NG, et al. Macrolide-resistance determination and molecular typing of *Mycoplasma pneumoniae* in respiratory specimens collected between 1997 and 2008 in The Netherlands. J Clin Microbiol 2012;50:1999–2004. <https://doi.org/10.1128/JCM.00400-12>.
- [12] Peuchant O, Menard A, Renaudin H, Morozumi M, Ubukata K, Bebear C, et al. Increased macrolide resistance of *Mycoplasma pneumoniae* in France directly detected in clinical specimens by real-time PCR and melting curve analysis. J Antimicrob Chemother 2009;64:52–8.
- [13] Chironna M, Sallustio A, Esposito S, Perulli M, Chinellato I, Di Bari C, et al. Emergence of macrolide-resistant strains during an outbreak of *Mycoplasma pneumoniae* infections in children. J Antimicrob Chemother 2011;66:734–7.
- [14] Achermann R, Suter K, Kronenberg A, Gyger P, Mühlemann K, Zimmerli W, et al. Antibiotic use in adult outpatients in Switzerland in relation to regions, seasonality and point of care tests. Clin Microbiol Infect 2011;17:855–61.